

## VU Research Portal

### Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene

Renssen, H.; Goosse, H.; Fichefet, T.

***published in***

Geophysical Research Letters  
2005

***DOI (link to publisher)***

[10.1029/2005GL022462](https://doi.org/10.1029/2005GL022462)

***document version***

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

***citation for published version (APA)***

Renssen, H., Goosse, H., & Fichefet, T. (2005). Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene. *Geophysical Research Letters*, 32(L08711).  
<https://doi.org/10.1029/2005GL022462>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene

Hans Renssen

Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

Hugues Goosse and Thierry Fichefet

Institut d'Astronomie et de Géophysique Georges Lemaître, Université Catholique de Louvain, Louvain-la-Neuve, Belgium

Received 17 January 2005; revised 15 March 2005; accepted 4 April 2005; published 30 April 2005.

[1] The Holocene North Atlantic deep-water formation is studied in a 9,000-year long simulation with a coupled climate model of intermediate complexity, forced by changes in orbital forcing and atmospheric trace gas concentrations. During the experiment, deep-water formation in the Nordic Seas is reduced due to an enhanced influx of sea ice from the Central Arctic, decreasing both surface salinity and density, whereas deep-water formation in the Labrador Sea increases due to surface cooling. This leads to changes in the distribution of oceanic heat transported northwards by the Atlantic Ocean, with less heat released ( $-120 \text{ Wm}^{-2}$  in February) in the Nordic Seas, amplifying the surface cooling and increasing the sea-ice cover. In the Labrador Sea, the oceanic heat release increases slightly ( $+14 \text{ Wm}^{-2}$ ), thus dampening locally the cooling trend. The overall Atlantic overturning strength remains constant throughout the experiment. Over the Nordic Seas, reduced evaporation contributes to the surface freshening. **Citation:** Renssen, H., H. Goosse, and T. Fichefet (2005), Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene, *Geophys. Res. Lett.*, 32, L08711, doi:10.1029/2005GL022462.

## 1. Introduction

[2] In the present-day North Atlantic Ocean the two main sites of deep-water formation are the Nordic Seas and the Labrador Sea [Weaver *et al.*, 1999]. In the Nordic Seas, Northeast Atlantic Deep Water (NEADW) is formed, which flows across the Greenland-Scotland ridge to fill the deep levels of the Atlantic, while in the Labrador Sea formation of intermediate water takes place that is known as Labrador Sea Water (LSW). The rate of deep-water formation is an important constituent of the total overturning circulation and the associated northward heat transport in the Atlantic Ocean. There is a growing concern that the North Atlantic Deep-Water formation might be reduced in a warmer climate [e.g., Wood *et al.*, 1999; Hansen *et al.*, 2004]. It is therefore important to understand the natural long-term variations in deep-water formation.

[3] Proxy studies suggest that deep-water formation in the Nordic Seas decreased after 7k (thousand year before


present [e.g., Rasmussen *et al.*, 2002]). The opposite trend has been found in the Labrador Sea, where data indicate that a modern-like circulation started slightly before 7k and strengthened afterwards [Hillaire-Marcel *et al.*, 2001; Solignac *et al.*, 2004]. Recent model studies have not been able to capture these contrasting trends [Cottet-Puinel *et al.*, 2004], leaving the underlying mechanism poorly understood. Therefore, we study here the Holocene evolution of deep-water formation in a 9,000-year long transient simulation performed with a coupled global climate model of intermediate complexity.

## 2. Experimental Design

[4] We applied version 3 of ECBilt-CLIO-VECODE, which describes the coupled atmosphere-sea ice-ocean-vegetation system in three dimensions. The atmospheric component is ECBilt, a quasi-geostrophic model with 3 vertical layers and T21 horizontal resolution [Opsteegh *et al.*, 1998]. The oceanic component CLIO consists of a free-surface, primitive-equation oceanic general circulation model coupled to a dynamic-thermodynamic sea-ice model [Goosse and Fichefet, 1999]. CLIO includes 20 levels in the vertical and has a  $3^\circ \times 3^\circ$  latitude-longitude horizontal resolution. VECODE is a model that simulates the dynamics of two main terrestrial plant functional types, trees and grasses, and desert as a dummy type [Brovkin *et al.*, 2002]. Details about the model are available at <http://www.knmi.nl/onderzk/CKO/ecbilt.html>.

[5] To simulate the long-term Holocene climate evolution, we performed a 9,000-year long experiment forced by changes in orbital parameters [Berger, 1978] and concentrations in atmospheric  $\text{CO}_2$  and  $\text{CH}_4$  [Raynaud *et al.*, 2000]. All other forcings (i.e., solar constant, land-sea-ice mask) were kept constant at preindustrial values. The initial conditions for the experiment were derived from a simulation that was run until equilibrium with 9k insolation and trace gas concentrations.

[6] In an earlier paper, we showed that the temperature and precipitation changes north of  $60^\circ\text{N}$  were in general agreement with proxy evidence [Renssen *et al.*, 2005]. The simulated response to orbital and greenhouse gas forcing experienced an early optimum (9–8 k) in most of the Arctic, followed by a 1 to  $3^\circ\text{C}$  decrease in mean annual temperatures, a reduction in summer precipitation and an expansion of sea-ice cover. Here we focus on the changes in the Labrador Sea and Nordic Seas in winter, as this is the



**Figure 1.** Simulated evolution of key parameters in the Nordic Seas (black line) and Labrador Sea (grey line) during February (i.e., month with maximum convection in the Labrador Sea). Results are averaged over 6 grid cells centered on the main convection sites. The lines depict the 100-point running means. (a) SST, (b) SSS, (c) surface density (note that  $1000 \text{ kg m}^{-3}$  has been subtracted for convenience) and (d) convection depth (note that maximum convection depths are 2060 m and 1215 m for Nordic and Labrador Seas, respectively).

season during which deep water is primarily formed under influence of the strong surface cooling.

### 3. Results and Discussion

[7] In the Labrador Sea and Nordic Seas, the winter sea surface salinity (SSS) and temperature (SST) are decreasing during the course of the experiment, leading to fresher and cooler surface waters (Figures 1a–1d and Table 1). These trends have opposite effects on the surface water density, as cooling leads to an increase in density, while freshening results in a decreased density. In the Nordic Seas the changes are relatively large. Compared to the Labrador

Sea, the cooling in the Nordic Sea is twice as large ( $-1.4^\circ\text{C}$  vs.  $-0.7^\circ\text{C}$ ), while the decrease in SSS is five times stronger ( $-0.26 \text{ psu}$  vs.  $-0.05 \text{ psu}$ ). These simulated trends are consistent with proxy-based reconstructions from the Labrador Sea [Solignac *et al.*, 2004] and the Nordic Seas [Koç *et al.*, 1993]. In the Labrador Sea, the very small 9k to 0k surface freshening of  $0.05 \text{ psu}$  is overwhelmed by the surface cooling, resulting in denser surface waters (Figure 1c) and deeper convection ( $+145 \text{ m}$  from 9k to 0k; Figures 1d and 2a). In the Nordic Seas, on the other hand, the long-term freshening is more substantial ( $-0.26 \text{ psu}$ ), leading to a reduced surface density (Figure 1c) and convection depth ( $-381 \text{ m}$  from 9k to 0k; Figures 1d and 2a), despite the  $1.4^\circ\text{C}$  ocean surface cooling. We found similar contrasting trends in deep-water formation between the Labrador Sea and Nordic Seas in sensitivity experiments with a slightly different setup than our main simulation [see, e.g., Renssen *et al.*, 2005], suggesting that this is a robust result in our model under Holocene forcings.

[8] The long-term winter cooling trends in both convection regions can be linked to changes in insolation. As explained by Renssen *et al.* [2005], the temperature over the Arctic lags the orbital forcing by several months due to the thermal inertia of the system, so that the early Holocene winters were warmer than today throughout the Arctic, despite the reduced insolation during winter. In addition to orbital forcing, the temperatures over the convection regions are influenced by processes that operate on a more local scale.

[9] In the Nordic Seas, the noted decrease in deep-water formation is accompanied by a 9k-to-0k reduction in *local* overturning of  $0.4 \text{ Sv}$  ( $1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$ ) or 12% on an annual basis [Renssen *et al.*, 2005]. This reduction does not influence the overall overturning rate of the Atlantic Ocean, as the weaker flow in the Nordic Seas is compensated by the enhanced contribution of LSW. As a result, there is even a slight (statistically insignificant) increase in the total Atlantic deep-water export at  $20^\circ\text{S}$  (from  $13.5 \text{ Sv}$  at 9k to  $13.8 \text{ Sv}$  at 0k) and in total deep-water production (from  $26.3 \text{ Sv}$  at 9k to  $27.4 \text{ Sv}$  at 0k). The total northward heat transport in the Atlantic Ocean remains at a constant level ( $0.32 \times 10^{15} \text{ W}$  at  $30^\circ\text{S}$ ) from 9k to 0k. These model results agree with available proxy records covering the Holocene, which suggest a weakening of the NEADW production [e.g., Solignac *et al.*, 2004], while showing no trend in the long-term overall Atlantic overturning circulation [e.g., Oppo *et al.*, 2003]. The changes in deep-water formation and local overturning circulation have an impact on the *distribution* of the heat transported northward by the

**Table 1.** Simulated 9k to 0k Changes in February, Averaged Over the Two Convection Regions

	Labrador Sea	Nordic Seas
SST ( $^\circ\text{C}$ )	−0.7	−1.4
SAT ( $^\circ\text{C}$ )	−0.7	−10.5
SSS (psu)	−0.05	−0.26
Surface density ( $\text{kg m}^{-3}$ )	+0.04	−0.11
Convection depth (m)	−145	+381
Ocean-to-atmosphere heat flux ( $\text{W m}^{-2}$ )	+14	−120
Sea-ice cover (%)	0	+27
Precipitation (mm)	−3	−16
Evaporation (mm)	0	−41





Svalbard region [see *Goosse et al.*, 2003; *Goosse and Renssen*, 2004].

[14] It should be noted that we have not included the effect of melting ice sheets in our experiment. Reconstructions indicate that the last remnants of the Laurentide Ice sheet melted around 7k [Peltier, 1994]. The associated meltwater flux is estimated at 0.08 Sv between 9 and 8k, of which only a portion drained directly into the Labrador Sea [Licciardi *et al.*, 1999]. After 8k the meltwater flux from the Laurentide ice sheet became very small (less than 0.01 Sv). Consequently, in the first thousand years of our experiment, the SSS in the Labrador Sea is probably overestimated, implying that the simulated upward trend in deep-water formation was more pronounced than in our simulation. This would be consistent with reconstructions that suggest that in the Labrador Sea a modern-like circulation started around 7k [Hillaire-Marcel *et al.*, 2001; Solignac *et al.*, 2004] and with the model simulations of Cottet-Puinel *et al.* [2004] that overestimated LSW formation when Laurentide meltwater was not accounted for.

#### 4. Concluding Remarks

[15] Our simulations suggest that in the warmer early Holocene climate, deep-water formation was stronger in the Nordic Seas than today, primarily because of a reduced sea-ice cover in the early Holocene and a relatively strong evaporation. In contrast, the opposite trend is found for the Labrador Sea, since here the simulated deep-water formation is governed by the surface cooling, which is less pronounced in the early Holocene than in the modern climate. These simulation results are in good agreement with available proxy evidence, suggesting that our findings provide a reasonable explanation for the observed Holocene trends in North Atlantic deep-water formation.

[16] **Acknowledgments.** HR is supported by the Netherlands Organization for Scientific Research (NWO). HG is Research Associate at the Belgian National Fund for Scientific Research. This study is part of the Belgian Second Multiannual Scientific Support Plan for a Sustainable Development Policy (Belgian Federal Science Policy Office, contract EV/10/9A and EV/10/7D) and the European Research Programme on Environment and Sustainable Development (contracts EVK2-2001-00263 and EVK2-CT-2002-00153).

#### References

- Berger, A. L. (1978), Long-term variations of daily insolation and Quaternary climatic changes, *J. Atmos. Sci.*, **35**, 2363–2367.
- Brovkin, V., J. Bendtsen, M. Claussen, A. Ganopolski, C. Kubatzki, V. Petoukhov, and A. Andreev (2002), Carbon cycle, vegetation, and climate dynamics in the Holocene: Experiments with the CLIMBER-2 model, *Global Biogeochem. Cycles*, **16**(4), 1139, doi:10.1029/2001GB001662.
- Cottet-Puinel, M., A. J. Weaver, C. Hillaire-Marcel, A. de Vernal, P. U. Clark, and M. Eby (2004), Variation of Labrador Sea Water formation over the last glacial cycle in a climate model of intermediate complexity, *Quat. Sci. Rev.*, **23**, 449–465.
- Goosse, H., and T. Fichefet (1999), Importance of ice-ocean interactions for the global ocean circulation: A model study, *J. Geophys. Res.*, **104**, 23,337–23,355.
- Goosse, H., and H. Renssen (2004), Exciting natural modes of variability by solar and volcanic forcing: Idealized and realistic experiments, *Clim. Dyn.*, **23**, 153–163.
- Goosse, H., F. M. Selten, R. J. Haarsma, and J. D. Opsteegh (2003), Large sea-ice volume anomalies simulated in a coupled climate model, *Clim. Dyn.*, **20**, 523–536.
- Hansen, B., S. Osterhus, D. Quadfasel, and W. Turrell (2004), Already the day after tomorrow?, *Science*, **305**, 953–954.
- Hillaire-Marcel, C., A. de Vernal, G. Bilodeau, and A. J. Weaver (2001), Absence of deep-water formation in the Labrador Sea during the last interglacial period, *Nature*, **410**, 1073–1077.
- Koç, N., E. Jansen, and H. Hafliðason (1993), Paleooceanographic reconstructions of surface ocean conditions in the Greenland, Iceland and Norwegian Seas through the last 14 ka based on diatoms, *Quat. Sci. Rev.*, **12**, 115–140.
- Licciardi, J. M., J. T. Teller, and P. U. Clark (1999), Freshwater routing by the Laurentide ice sheet during the last deglaciation, in *Mechanisms of Global Climate Change at Millennial Time Scale*, *Geophys. Monogr. Ser.*, vol. 112, edited by P. U. Clark, R. S. Webb, and L. D. Keigwin, pp. 177–201, AGU, Washington, D. C.
- Oppo, D. W., J. F. McManus, and J. L. Cullen (2003), Deepwater variability in the Holocene epoch, *Nature*, **422**, 277–278.
- Opsteegh, J. D., R. J. Haarsma, F. M. Selten, and A. Kattenberg (1998), ECBILT: A dynamic alternative to mixed boundary conditions in ocean models, *Tellus, Ser. A*, **50**, 348–367.
- Peltier, W. R. (1994), Ice age paleotopography, *Science*, **265**, 195–201.
- Rasmussen, T. L., D. Bäckström, J. Heinemeier, D. Klitgaard-Kristensen, P. C. Knutz, A. Kuijpers, S. Lassen, E. Thomsen, S. R. Troelstra, and T. C. E. van Weering (2002), The Faroe-Shetland Gateway: Late Quaternary water mass exchange between the Nordic seas and the north-eastern Atlantic, *Mar. Geol.*, **188**, 165–192.
- Raynaud, D., J.-M. Barnola, J. Chappellaz, T. Blunier, A. Indermühle, and B. Stauffer (2000), The ice record of greenhouse gases: A view in the context of future changes, *Quat. Sci. Rev.*, **19**, 9–17.
- Renssen, H., H. Goosse, T. Fichefet, V. Brovkin, E. Driesschaert, and F. Wolk (2005), Simulating the Holocene climate evolution at northern high latitudes using a coupled atmosphere-sea ice-ocean-vegetation model, *Clim. Dyn.*, **24**, 23–43.
- Solignac, S., A. de Vernal, and C. Hillaire-Marcel (2004), Holocene sea-surface conditions in the North Atlantic—Contrasted trends and regimes in the western and eastern sectors (Labrador Sea vs. Iceland Basin), *Quat. Sci. Rev.*, **23**, 319–334.
- Weaver, A. J., C. M. Bitz, A. F. Fanning, and M. M. Holland (1999), Thermohaline circulation: High-latitude phenomena and the difference between Pacific and Atlantic, *Annu. Rev. Earth Planet. Sci.*, **27**, 231–285.
- Wood, R. A., A. B. Keen, J. F. B. Mitchell, and J. M. Gregory (1999), Changing spatial structure of the thermohaline circulation in response to atmospheric CO<sub>2</sub> forcing in a climate model, *Nature*, **399**, 572–575.

T. Fichefet and H. Goosse, Institut d'Astronomie et de Géophysique Georges Lemaître, Université Catholique de Louvain, 2, chemin du Cyclotron, B-1348, Louvain-la-Neuve, Belgium.

H. Renssen, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, NL-108HV, Amsterdam, Netherlands. (hans.rensSEN@geo.falw.vu.nl)